



Geochemical Studies of Ground Water at the Eastern Side Dindigul Rock Fort due to the Impact of Sewage and Industrial Effluent

A. Pandia Rajan^{1*}, M. Velusamy², R. Sukumar³, I. N. Karthika⁴, M. S. Dheenadhayalan^{5*}

^{1*,2,3,5*}Department of Chemistry, GTN Arts College, Dindigul, TN, India.

⁴P.S.N.A College of Engineering and Technology, Dindigul, TN, India.

Received : 12.04.2014 Accepted : 29.05.2014

Abstract

An attempt has been made to study the impact of untreated sewage water and industrial effluent in the ground water sources from the houses located at the foot hills of Dindigul Rock fort. The pond has become the collection of sewage water from Dindigul town. About 10 ponds in Dindigul town discharges through various canals and finally reaches the Rock fort pond behind Taluk Office. Hence the water in the pond as well as the ground water sources around the various residential area like Kamarajapuram, due to high degree of pollution many sources has to be abandoned ground water analysis at four different sites at four direction reveals that the water quality parameters are higher than the permitted level. As per CPHEEO standard specifically high turbidity, high TDS and higher electrical conductivity values indicates that the water cannot be used for human consumption or any other use. The adjoining ground water sources are mostly affected and the water becomes very salty with very high TDS. Hence the polluted water is suggested to water treatment using Reverse Osmosis System.

Keywords: CPHEEO; Ground Water; Industry effluent; Sewage.

1. INTRODUCTION

1.1 Water Pollution

Water pollution can be defined in many ways. Usually, it means one or more substances have built up in water to such an extent that they cause problems for animals or people. Oceans, lakes, rivers, and other inland waters can naturally clean up a certain amount of pollution by dispersing it harmlessly. If you poured a cup of black ink into a river, the ink would quickly disappear into the river's much larger volume of clean water. The ink would still be there in the river, but in

such a low concentration that you would not be able to see it. At such low levels, the chemicals in the ink probably would not present any real problem. However, if you poured gallons of ink into a river every few seconds through a pipe, the river would quickly turn black. The chemicals in the ink could very quickly have an effect on the quality of the water. This, in turn, could affect the health of all the plants, animals, and humans whose lives depend on the river.

1.2 Sewage

With billions of people on the planet, disposing of sewage waste is a major problem. According to 2004 figures from the World Health Organization, some 1.1 billion people (16 percent of the world's population)

*A. Pandia Rajan Tel. no.: 9380939942
E-mail: chempandian@gmail.com

don't have access to safe drinking water, while 2.6 billion (40 percent of the world's population) don't have proper sanitation (hygienic toilet facilities); the position hasn't improved much since. Sewage disposal affects people's immediate environments and leads to water-related illnesses such as diarrhea that kills 3-4 million children each year. (According to the World Health Organization, water-related diseases could kill 135 million people by 2020.) In developed countries, most people have flush toilets that take sewage waste quickly and hygienically away from their homes.

Effluent irrigation has been practiced for centuries throughout the world (Shuval *et al.* 1986; Tripathi *et al.* 2011). It provides farmers with a nutrient enriched water supply and society with a reliable and inexpensive system for wastewater treatment and disposal (Feigin *et al.* 1991). In India also being a cheap source of irrigation. Rapid industrialization, population explosion and more urbanization in India have created enormous problems of environmental pollution in terms of generating the variable quantity and quality of solid and liquid wastes. In developing countries, there has not been much emphasis on the installation of sewage treatment plants and all the industrial effluents are generally discharged in to the sewage system. The sewage waters are used as potential source irrigation for raising vegetables and fodder crops around the sewage disposal sites which are directly or indirectly consumed by human beings.

Soil contamination by sewage and industrial effluents has affected adversely both soil health and crop productivity. Sewage and industrial effluents are the rich sources of both beneficial as well as harmful elements. Since some of these effluents are a rich source of plant nutrients, therefore soil provides the logical sink for their disposal. But many untreated and contaminated sewage and industrial effluents may have high concentration of several heavy metals such as Cd, Ni, Pb and Cr (Arora *et al.* 1985). Their continuous disposal on agricultural soils has resulted in soil sickness (Narwal *et al.* 1988) and accumulation of some

of the toxic metals in soil (Adhikari *et al.* 1993; Antil, 2005; Gupta *et al.* 2002, 1998; Kharche *et al.* 2011) which may pose serious human and animal health. Ground water in Punjab has been contaminated by Hg and Pb to such an extent that it is causing DNA of the people, who drink it, to mutate (Bajwa, 2008). The present chapter, therefore, discusses the composition of sewage and industrial effluents in India and their possible effect on soil-plant health.

Over two thirds of Earth's surface is covered by water; less than a third is taken up by land. As Earth's population continues to grow, people are putting ever-increasing pressure on the planet's water resources. In a sense, our oceans, rivers, and other inland waters are being "squeezed" by human activities—not so they take up less room, but so their quality is reduced. Poorer water quality means water pollution.

2. SCOPE AND OBJECTIVES OF THE STUDY

The volume of sewage water discharged from the house in dindigul town through nine ponds are increasing day by day finally the sewage water is collected at the foot of the Rock fort in a pond, similarly the industrial waste water from dyeing units and other small scale industries. The sewage water and the industries effluent from the canals discharged the polluted water in to the pond without any treatment. The option of treatment plant to treat the sewage water and the effluent may lead to spoilage of environment one time morning people will not be able to get good quality of drinking water from the ground water in and around the Dindigul Rock fort area. The environmental damage caused by water pollution by the discharge of sewage water and industrial effluent in the pond at foot hills of Dindigul Rock fort.

2.1 Objectives

- To evaluate the physic chemical parameters of the water in the pond at the foot hills of Dindigul Rock fort.

- To evaluate the physico-chemical parameters of the groundwater present in the well and bore wells around the pond located at the foot hills of Dindigul Rock fort.
- To suggest a suitable remedial measure to treat the groundwater using RO system.

3. MATERIALS & METHODS

The pond located at the foot hills of Dindigul Rock fort has become the collection of sewage water and industrial effluent from the Dindigul town due to slope and gradient all the sewage water from 10 ponds in Dindigul town reaches the rock fort pond on the Northern side of Dindigul Rock fort. Hence the water in the pond as well as the groundwater sources in and around the Rock fort area at a radius of 2 km are completely polluted due to the continuous discharge and percolations of the sewage water to the groundwater.

The absence of any organized sewage and industrial effluent systems and the sewage treatment plant in Dindigul town, the community wastes often find their way to the nearby lake, pond and river. During the rainy seasons almost all the 10 ponds located in Dindigul at different areas in all four directions discharge the sewage water through big canals and finally reach the pond at the foot hills of the Rock fort which leads to groundwater pollution at a radius of 5 km around the Rock fort. The rainwater which percolates into the ground is the main source of water to the residential areas like Kamarajapuram, Sannieswaran kovil and Sreenivasaperumal kovil located at the bottom of the Rock fort are very much affected. During the survey people in the residential area are not using the groundwater due to its high salty taste with high turbidity.

3.1 Analysis of the Contaminated Water

S.No	Parameter	Method Of Analysis
1	Colour	Visual comparison
2	Turbidity	Neplo turbidity meter
3	TDS	Conductivity method
4	Electrical conductivity	Conductivity meter
5	pH	pH Meter
6	Total hardness	EDTA Titrimetric method
7	Calcium	EDTA Titrimetric method
8	Magnesium	Calculation from Total Hardness
9	Iron	Spectrophotometer
10	Ammonia	Nessler's Method
11	Nitrite	Spectrophotometer
12	Nitrate	Spectrophotometer
13	Chloride	Silver nitrate
14	Fluoride	Colorimetric meter
15	Sulphate	Turbidity method
16	Phosphate	Spectrophotometer

4. RESULTS & DISCUSSIONS

4.1 Drinking Water Standards

Raw water quality and standards depend upon the end use. The four main uses are municipal, industrial, agricultural and recreational (fish and wildlife). As water quality is degraded day by day, so it becomes very important to set the drinking water standards for the safety of water of our limited resources. Different agencies have set environmental standards for safe drinking water like Bureau of Indian Standards (BIS), World Health Organization (WHO), and European Economic Community (EEC) etc. Drinking water standards are regulations that the Bureau of Indian Standards (BIS) set to control the level of contamination in the drinking water. The Bureau of Indian Standards considers the inputs from many organizations i.e. Central, State,

4.1. Comprehensive Table Water Quality Analysis

Sample collection	BIS Limit	S1	S2	S3	S4
Appearance		Turbid	Clear	Clear	Clear
colour(Pt.Co-Scale)	<u>5</u>	Blackish	Colourless	Colourless	Colourless
Turbidity NT units	5	60	7	6	3
Total dissolved solids mg/L	<u>500</u>	1409	808	904	653
Electrical conductivity in Micro mhos/cm		2072	1189	1329	960
pH	7.0-8.5	7.49	7.69	8.03	7.63
Total hardness as CaCO ₃ (mg/L)	300	700	268	340	272
Calcium as Ca mg/L	75	144	64	74	61
Magnesium as Mg mg/L	30	82	26	37	29
Sodium as Na	-	120	124	112	78
Potassium as K	-	30	32	28	20
Iron as Fe mg/L	<u>0.3</u>	4.31	1.02	0.59	0.86
Ammonia as NH ₃ mg/L	-	5.34	1.09	0.76	0.59
Nitrite as NO ₂ mg/L	-	0.73	0.33	0.22	0.16
Nitrate as NO ₃ mg/L	<u>45</u>	16	13	6	5
Chloride as Cl mg/L	250	440	176	224	162
Fluoride as F mg/L	1	1.2	0.4	0.8	0.4
Sulphate as SO ₄ mg/L	200	117	116	76	62
Phosphate as PO ₄ mg/L	-	2.50	0.90	0.72	0.63

Semi Government, Municipal Corporation, Public Health Organization, etc., throughout the standard setting process.

4.2. Sensitive Parameters

Parameters like TDS, EC, hardness, calcium, magnesium, Iron, fluoride and Sulphate are taken as

sensitive parameters to indicate the water pollution by industrial effluent from various sources. It is observed that the values are higher compared the BIS Standards.

5. CONCLUSION

An attempt has been made to study the impact of untreated sewage water and industrial effluent in the

ground water sources from the houses located at the foot hills of Dindigul Rock fort. The pond has become the collection of sewage water from Dindigul town About 10 ponds in Dindigul town discharges through various canals and finally reaches the Rock fort pond behind Taluk Office. Hence the water in the pond as well as the ground water sources around the various residential area like Kamarajapuram, due to high degree of pollution many sources has to be abandoned ground water analysis at four different sites at four direction reveals that the water quality parameters are higher than the permitted level. As per CPHEEO standard specifically high turbidity high TDS and higher Electrical conductivity values indicates that the water cannot be used for human consumption or any other use. People have to depend only on municipal water sources from Authoor, For other uses the ground water can be treated using RO system. The adjoining ground water sources are mostly affected and the water becomes very salty with very high TDS. Hence the polluted water is suggested to water treatment using Reverse Osmosis System.

ACKNOWLEDGEMENT

The authors are thankful to their concerned TWAD board and the Management of the college for constant encouragement and financial support to promote such innovative solution for the environmental pollution problem.

REFERENCES

- Adhikari, S., Gupta, S. K. and Banerjee, S. K., Heavy metals content of city sewage and sludge, *Journal of Indian Society of Soil Science*, 41, 160-172(1993).
- Antil, R. S. and Narwal, R. P., Problems and prospectus of utilization of sewer water in Haryana. In: Management of organic wastes for crop production, Kapoor, K. K, Sharma, P. K., Dudeja, S. S. and Kundu, B. S., (Ed.), 159-168, Department of Microbiology, CCS Haryana Agricultural University, Hisar, India(2005).
- Arora, B. R., Azad, A. S., Singh, B. and Sekhon, G. S., Pollution potential of municipal waste waters of Ludhiana, Punjab. *Indian Journal of Ecology*, 12, 1-7(1985).
- Bajwa, H., Water contamination causing DNA mutation. *The Indian Express*, 5, Feb. 16, 2008, Chandigarh edition, India(2008).
- Feigin, A., Ravina, I. and Shalhevet, J., Irrigation with treated sewage effluent. Management for Environmental Protection. Advanced series in Agricultural Sciences 17, 224, Springer-Verla(1991).
- Gupta, S. K. and Mitra, A., In: Advances in land resource management for 21st century, Soil conservation society of India, New Delhi, 446-469(2002).
[doi:10.1016/S0960-8524\(98\)00003-0](https://doi.org/10.1016/S0960-8524(98)00003-0)
- Gupta, A. P., Narwal, R. P. and Antil, R. S., Sewer waters composition and their effect on soil properties. *Bioresour. Technol.*, 65, 171-173(1998).
- Kharche, V. K., Desai, V. N. and Pharande, A. L., Effect of sewage irrigation on soil properties, essential nutrients and pollutant element status of soils and plants in a vegetable growing area around Ahmednagar city in Maharashtra, *Journal of Indian Society of Soil Science*, 59, 177-184(2011).
- Narwal, R. P., Singh, M. and Gupta, A. P., Effect of different sources of irrigation on the physico-chemical properties of soil, *Indian Journal of Environment and Agriculture*, 3, 27-34(1988).
- Shuval, H. I., Adin, A., Fattal, B., Rawitz, E. and Yekutieli, P., Wastewater irrigation in developing countries, *Health effects and technical solutions*, 325, World Bank Technical Report 51(1986).
- Tripathi, D. M., Tripathi, S. and Tripathi, B. D., Implications of secondary treated distillery effluent irrigation on soil cellulase and urease activities, *Journal of Environment Protection*, 2: 655-661(2011).
[doi:10.4236/jep.2011.25075](https://doi.org/10.4236/jep.2011.25075)